



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Long-term effect of yolk carotenoid levels on testis size in a precocial bird

Giraudeau, Mathieu ; Ziegler, Ann-Kathrin ; Tschirren, Barbara

Abstract: Conditions experienced during prenatal development can have long-lasting organizational effects on offspring. Maternal carotenoids deposited in the eggs of birds and other oviparous species play an important role during fast embryonic growth and chick development through their antioxidant properties. However, the long-term consequences of variation in maternal carotenoid transfer for the offspring have seldom been considered. Since plasma carotenoid levels at adulthood are known to influence testis size and yolk carotenoid levels influence the ability to extract carotenoids later in life, we hypothesized that maternally transmitted carotenoids might influence gonad size at adulthood. Here, we showed that male Japanese quail (*Coturnix japonica*) originating from a carotenoid-enriched egg had smaller testes than control individuals at adulthood. This result shows that yolk carotenoids have long-term organizational effects. In addition, given that carotenoid intake at sexual maturity increases sperm quality and that a decreased testis size is associated with a lower sperm production, we propose that carotenoid exposure during embryo development might influence a trade-off between ejaculate size and sperm quality.

DOI: <https://doi.org/10.1098/rsbl.2016.0008>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-126756>

Journal Article

Accepted Version

Originally published at:

Giraudeau, Mathieu; Ziegler, Ann-Kathrin; Tschirren, Barbara (2016). Long-term effect of yolk carotenoid levels on testis size in a precocial bird. *Biology Letters*, 12(4):20160008.

DOI: <https://doi.org/10.1098/rsbl.2016.0008>

**LONG-TERM EFFECT OF YOLK CAROTENOID LEVELS ON TESTIS SIZE IN A
PRECOCIAL BIRD**

GIRAUDEAU MATHIEU^{*,1,2,3}, ZIEGLER ANN-KATHRIN¹ AND TSCHIRREN
BARBARA¹

¹ Department of Evolutionary Biology and Environmental Studies, University of Zurich,
Switzerland.

² Arizona State University, School of Life Sciences, Tempe, AZ, USA.

³ Centre for Ecology & Conservation, College of Life and Environmental Sciences, University
of Exeter, Penryn, UK.

*E-mail: giraudeau.mathieu@gmail.com

Keywords: carotenoids, maternal effects, long-term effects, testis size

13 pages

2500 words

1 figure, 0 table

ABSTRACT

Conditions experienced during prenatal development can have long-lasting organizational effects on offspring. Maternal carotenoids deposited in the eggs of birds and other oviparous species play an important role during fast embryonic growth and chick development through their antioxidant properties. However, the long-term consequences of variation in maternal carotenoid transfer for the offspring have been seldom considered. Since plasma carotenoid levels at adulthood are known to influence testis size and yolk carotenoid levels influence the ability to extract carotenoids later in life, we hypothesized that maternally transmitted carotenoids might influence gonad size at adulthood. Here, we showed that male Japanese quail (*Coturnix japonica*) originating from a carotenoid-enriched egg had smaller testes than control individuals at adulthood. This result shows that yolk carotenoids have long-term organizational effects. In addition, given that carotenoid intake at sexual maturity increases sperm quality and that a decreased testis size is associated with a lower sperm production, we propose that carotenoid exposure during embryo development might influence the trade-off between ejaculate size and sperm quality.

INTRODUCTION

Conditions experienced during embryo development can have major organizational effects that can last until adulthood [1]. However, although numerous studies have documented short-term effects of prenatal conditions on offspring phenotype [2], the long-term consequences for fitness-related traits are still poorly understood.

Maternally transmitted antioxidants might influence embryonic developmental trajectory due to their capacity to scavenge the reactive oxygen species produced during development and the challenging period of rapid growth [3]. Several studies have examined the importance of these compounds during development using dietary carotenoid supplementation of laying females to indirectly manipulate yolk antioxidants levels (i.e. carotenoids) and have shown effects of this treatment on nestlings' condition or carotenoid levels just a few weeks after hatching [4,5,6]. However, these results must be considered cautiously since these effects may have been mediated by other effects of these dietary manipulations on maternal physiology and/or the differential allocation of other maternally transmitted compounds.

So far, only two studies have directly manipulated yolk carotenoid concentrations through *in ovo* injections, showing that these maternally transmitted compounds have the potential to influence chick growth, immunocompetence and antioxidation capacity [7,8,9]. The potential long-term effects of prenatal antioxidant exposure have so far only been examined in one study in which male barn swallows (*Hirundo rustica*) hatched from eggs injected with vitamin E arrived earlier at their breeding grounds than controls [10]. We thus need more studies where the long-term consequences of yolk antioxidant manipulations (through *in ovo* injections) are examined to determine the potential organizational effect of these maternally transmitted compounds.

Nutritional conditions during development and at adulthood have been shown to influence gonadal development in a variety of taxa (cockroach, [11]), humans [12], mallard [13]), but information on the importance of specific antioxidants (such as carotenoids) on gonadal growth is scarce. Carotenoids are present in both testes and seminal fluid and it has been proposed that these molecules might limit oxidative stress and allow optimal cell growth in testes [14], a tissue where the high rate of cell division might generate high levels of free radicals [15]. Given that variation in yolk carotenoid levels can influence the ability to extract or assimilate these compounds later in life [4] and that dietary carotenoid availability at adulthood influenced testes size [16], we hypothesized that maternally transmitted carotenoid may influence gonad size at adulthood. To test this hypothesis, we experimentally manipulated yolk lutein levels in Japanese quail eggs and measured the consequences of this treatment for the sons' testis size at adulthood.

METHODS

Unincubated Japanese quail (*Coturnix japonicus*) eggs were collected from 55 females of our captive breeding population and injected with either 15 µg of carotenoids (FloraGLO Lutein 20%, Kemin Foods, Des Moines, Iowa) dissolved in 15µL of safflower oil or with only safflower oil as a control. Lutein was chosen since it is the most abundant carotenoid found in Japanese quail eggs and the dose injected represents approximately one standard deviation of the yolk carotenoid content in this species [17]. The overall hatching success was 41.1% (Control = 38.5%, Carotenoid = 44.1%) and comparable to previous studies in Japanese quail [18, 19]. See ESM for details on the incubating and rearing conditions. One year post-hatch, 48 males were randomly selected and euthanized (29 control and 19 birds from carotenoid-injected eggs, from 40 females and 430 eggs injected ((226 control-injected and 204 carotenoid-

injected)). Both testes were collected and weighed to the nearest 1 mg. Tarsus length was measured to the nearest 0.1 mm.

Statistics

Testis mass was significantly repeatable between the left and right sides ($F_{1, 45} = 23.45$, $P < 0.001$), we thus used average values per bird for our analyses. In order to avoid pseudo-replication, we used the mean average testis mass of all male offspring per mother in our analyses since six mothers had more than one son. We ran general linear models to test whether the yolk carotenoid manipulation and bird size (tarsus length), as well as their interaction, affected testis size. All statistical analyses were run in R 3.01 (R Core Team 2013).

RESULTS

Bigger birds had bigger testes ($F_{1,37} = 7.77$, $P = 0.008$). Furthermore, testis size was significantly affected by the carotenoid treatment ($F_{1,37} = 4.52$, $P = 0.04$) with males originating from carotenoid-injected eggs having smaller testes than males originating from control eggs (figure 1). The interaction effect between carotenoid treatment and tarsus length was non-significant ($F_{1, 36} = 0.030$, $P = 0.86$). Yolk carotenoid treatment did not influence body size at adulthood ($F_{1, 38} = 1.41$, $P = 0.24$).

DISCUSSION

Here, we show that prenatal exposure to carotenoids has long-term effects on a primary sexual trait. Male Japanese quail originating from a carotenoid-injected egg had smaller testes than controls one year post-hatching. This result is in line with the idea that yolk carotenoids have organizational effects that last until adulthood, potentially due to their antioxidant

properties [20, but see 21] and / or their effect on gene expression, cell proliferation, and cell–
cell communication [22, 23].

Previous studies have indirectly manipulated yolk carotenoid levels through supplementation
of laying females and showed that chick absorption and utilization of carotenoids,
immunocompetence and plumage coloration were influenced by this dietary manipulation
several weeks after hatching [4,5,24]. Our result confirms these long-lasting effects of yolk
carotenoid levels using a direct *in ovo* manipulation of these maternally-transmitted compounds
and by measuring a primary sexual trait several months after the end of the developmental
phase. Given that only one carotenoid has been manipulated in our study, further work is needed
to investigate if also other yolk carotenoids have long-term effects on fitness-related traits.

We propose three non-mutually exclusive hypotheses that might explain the reduction
of testis size in males originating from carotenoid-enriched eggs. First, carotenoids may
influence the trade-off between ejaculate size and sperm quality [25]. Recent studies have
shown that an increased dietary intake of carotenoid leads to a reduction of testes size in
mallards (*Anas platyrhynchos*) [13] and that testis size and testis carotenoid concentrations are
negatively correlated in house finches (*Haemorrhous mexicanus*) [16]. Given that testis size is
positively associated with sperm production in birds [26], this strongly suggests that high
carotenoid intake decreases ejaculate size in birds. However, studies in various taxa have also
shown that an increased carotenoid intake at sexual maturity increases sperm quality [27, 28,
29], potentially through their antioxidant properties or the recycling of vitamin E molecules, a
major actor of spermatozoa protection from oxidative damage [29]. Thus, carotenoids seem to
stimulate the production of high-quality sperm, but in smaller amounts. Future studies should
test this hypothesis by simultaneously measuring ejaculate size and sperm quality in individuals

supplemented or not with carotenoid at different life stages (before or after hatching). Second, prenatal exposure to high levels of carotenoid may influence the trade-off between self-maintenance and survival toward a reduced reproductive investment during the first breeding event. However, this hypothesis seems unlikely since carotenoid supplementation has been shown to increase several components of reproductive investment in various species [30]. Finally, while we acknowledge the possibility that prenatal carotenoid exposure may have detrimental consequences for testes maturation and thus sperm production in our study, we believe that it is inconsistent with the accumulating evidence that carotenoid supplementation improves sperm quality [27, 29, but see 31]. In addition, the injected carotenoid dose was well within the natural range (Peluc *et al.* 2012) and yolk carotenoid levels after injection were not unnaturally high since females were fed with a low-carotenoid diet during the whole experiment.

To conclude, we show for the first time that yolk carotenoid levels have long-term effects on a primary sexual trait, strongly suggesting that maternally-transmitted antioxidants influence offspring fitness.

Acknowledgements: We thank Alison Pick and Pascale Hutter for help with data collection.

Ethics: All procedures conform to the relevant regulatory standards and were conducted under licences provided by the Veterinary Office of the Canton of Zurich, Switzerland (195/2010; 14/2014; 156).

Data accessibility: Data are available from the Dryad repository: <http://doi:10.5061/dryad.523t4>.

152 **Funding:** The study was supported by the Swiss National Science Foundation
153 (PP00P3_128386 and PP00P3_157455) and the Fonds zur Förderung des akademischen
154 Nachwuchses.


155 **Authors' contributions:** MG and AZ collected the data, MG and BT designed the study, BT
156 analyzed the data. MG and BT wrote the manuscript. All authors agree to be held accountable
157 for the content therein and gave final approval for publication.

158 **Competing interests:** We declare we have no competing interests.

159 Figure 1: Effect of yolk carotenoid injection on testis size one year post-hatching. Means \pm 1 SE
160 are shown.

161

162



REFERENCES

1. Mousseau TA, Fox CW. 1998 Maternal Effects as Adaptations. Oxford University Press, New York.
2. Groothuis TGG, Wendt M, von Engelhardt N, Carere C, Eising C. 2005 Maternal hormones as a tool to adjust offspring phenotype in avian species. *Neurosci. Biobehav. Rev.* **29**, 329–352.
3. Surai PF, Speake BK, Sparks NHC. 2001 Carotenoids in avian nutrition and embryonic development. 2. Antioxidant properties and discrimination in embryonic tissues. *J. Poultry Sci.* **38**, 117–145.
4. Koutsos EA, Clifford AJ, Calvert CC, Klasing KC. 2003 Maternal carotenoid status modifies the incorporation of dietary carotenoids into immune tissues of growing chickens (*Gallus gallus domesticus*). *J. Nutr.* **133**, 1132–1138.
5. McGraw KJ, Adkins-Regan E, Parker RS. 2005 Maternally derived carotenoid pigments affect offspring survival, sex ratio, and sexual attractiveness in a colorful songbird. *Naturwissenschaften* **92**, 375–380.
6. Grether GF, Kolluru GR, Lin K, Quiroz MA, Robertson G, Snyder AJ. 2008 Maternal effects of carotenoid consumption in guppies (*Poecilia reticulata*). *Funct. Ecol.* **22**, 294–302.
7. Saino N, Ferrari RP, Romano M, Martinelli R, Møller AP. 2003 Experimental manipulation of egg carotenoids affects immunity of barn swallow nestlings. *Proc R Soc Lond B Biol Sci.* **270**, 2485–2489.
8. Saino N, Romano M, Caprioli M, Rubolini D, Ambrosini R. 2011 Yolk carotenoids have sex-dependent effects on redox status and influence the resolution of growth trade-offs in yellow-legged gull chicks. *Behav. Ecol.* **22**, 411–421.

- 187 9. Romano M, Caprioli M, Ambrosini R, Rubolini D, Fasola M, Saino N. 2008 Maternal
188 allocation strategies and differential effects of yolk carotenoids on the phenotype and
189 viability of yellow-legged gull chicks in relation to sex and laying order. *J. Evol. Biol.*
190 **21**, 1826–1840.
- 191 10. Moller AP, Biard C, Karadas F, Rubolini D, Saino N, Surai PF. 2011 Maternal effects
192 and changing phenology of bird migration. *Climate research* **49**, 201-210.
- 193 11. Barrett ELB, Hunt J, Moore AJ, Moore PJ. 2009 Separate and combined effects of
194 nutrition during juvenile and sexual development on female life-history trajectories: the
195 thrifty phenotype in a cockroach. *Proc. R. Soc. B* **276**, 3257–3264.
- 196 12. Lummaa V. 2003 Early developmental conditions and reproductive success in humans:
197 downstream effects of prenatal famine, birthweight, and timing of birth. *Am. J. Hum.*
198 *Biol.* **15**, 370–379.
- 199 13. Butler MW, Karanfilian B, Homsher M, McGraw KJ. 2013 Carotenoid supplementation
200 during adulthood, but not development, decreases testis size in mallards. *Comparative*
201 *Biochemistry and Physiology, Part A* **166**, 465–469.
- 202 14. Blount JD, Møller AP, Houston DC. 2001 Antioxidants, showy males and sperm
203 quality. *Ecol. Letters* **4**, 393-396.
- 204 15. Taylor CT. 2001 Antioxidants and reactive oxygen species in human fertility. *Environ.*
205 *Tox. Pharm.* **10**, 189–198.
- 206 16. Rowe M, Tourville EA, McGraw KJ. 2012 Carotenoids in bird testes: links to body
207 carotenoid supplies, plumage coloration, bodymass and testes mass in house finches.
208 *Comp. Biochem. Physiol. B* **163**, 285–291.
- 209 17. Peluc SI, Reed WL, McGraw KJ, Gibbs P. 2012 Carotenoid supplementation and GnRH
210 challenges influence female endocrine physiology, immune function, and egg-yolk

- characteristics in Japanese quail (*Coturnix japonica*). *J. Comp. Physiol. B.* **182**, 687–702.
18. Daisley JN, Bromundt V, Mostl E, Kotrschal K. 2005 Enhanced yolk testosterone influences behavioral phenotype independent of sex in Japanese quail chicks *Coturnix japonica*. *Horm. Behav.* **47**, 185–194.
19. Hegyi G, Schwabl H. 2010 Do different yolk androgens exert similar effects on the morphology or behaviour of Japanese quail hatchlings *Coturnix japonica*? *J. Avian Biol.* **41**, 258–265.
20. Halliwell B, Gutteridge J. 2007 Free radicals in biology and medicine. Oxford: Oxford University Press.
21. Costantini D, Møller AP. 2008 Carotenoids are minor antioxidants for birds. *Funct Ecol.* **22**, 367–370.
22. Edge R, McGarvey DJ, Truscott TG. 1997 The carotenoids as antioxidants- a review. *J Photochem Photobiol B.* **41**, 189–200.
23. Chew BP, Park JS. 2004 Carotenoid action on the immune response. *J Nutr.* **134**, 257S–261S.
24. Biard C, Surai PF, Møller AP. 2007 An analysis of pre- and post-hatching maternal effects mediated by carotenoids in the blue tit. *J Evol Biol.* **20**, 326–339.
25. Immler S, Pitnick S, Parker GA, Durrant KL, Lüpold S, Calhim S, Birkhead T. 2011 Resolving variation in the reproductive tradeoff between sperm size and number. *PNAS* **108**, 5325–5330.
26. Møller AP. 1988 Testes size, ejaculate quality and sperm competition in birds. *Biol. J. Linn. Soc.* **33**, 273–283.

- 234 27. Helfenstein F, Losdat S, Møller AP, Blount JD, Richner H. 2010 Sperm of colourful
235 males are better protected against oxidative stress. *Ecol. Lett.* **13**, 213–222.
- 236 28. Taş M, Güney Saruhan B, Kurt D, Yokuş B, Denl M. 2010 Protective role of lycopene
237 on aflatoxin B1 induced changes sperm characteristics and testicular damages in rats.
238 Kafkas Univ. *Vet. Fak. Derg.* **16**, 597–604.
- 239 29. Almbro M, Dowling DK, Simmons LW. 2011 Effects of vitamin E and beta-carotene
240 on sperm competitiveness. *Ecol. Lett.* **14**, 891–5.
- 241 30. Blount JD, Houston DC, Surai PF, Møller AP. 2004 Egg-laying capacity is limited by
242 carotenoid pigment availability in wild gulls *Larus fuscus*. *Proc. R. Soc. B* **271**, S79–
243 S81.
- 244 31. Sullivan M, Brown AC, Clotfelter ED. 2014 Dietary carotenoids do not improve
245 motility or antioxidant capacity in cichlid fish sperm. *Fish Physiology and Biochemistry*
246 **40(5)**, 1399-1405.

ESM

Eggs were artificially incubated for 14 days at a temperature of 37.6°C and 55% humidity and then at 37.6°C and 80% humidity for the last 3 days. Chicks were reared in mixed treatment cohorts of 40 chicks for 2 weeks and in cohorts of 20 chicks for three more weeks. At the age of 5 weeks, chicks were released into outdoor aviaries. Adults and chicks received *ad libitum* water and commercial game bird mix low in carotenoid content during the whole experiment.